



Skin Des

Skin and Hand Disinfection

2-Propanol 70 % (V/V) lotion for hand disinfection, disinfection of skin before injections, for cold compression therapy

- bactericidal disinfectant (including TB)
- fungicidal
- limitedly virucidal in accordance with RKI recommendation (HBV, HIV, BVDV, HCV, vaccinia viruses, influenza viruses) and rotaviruses
- no spores due to filtering
- perfume-free

Product properties

Alcoholic lotion for hand and skin disinfection. Perfume-free and colourant-free. Particularly suitable for sensitive skin and skin tending to allergies.

Composition

Active ingredients in 100 g lotion: 2-Propanol 63,1 g; other ingredient: refined water.

Areas of Application

Hand disinfection and Surgical hand disinfection, disinfection of skin before simple injections and puncture of peripheral blood vessels, disinfection of skin before operations and punctures of joints, disinfection of sebaceous skin; cold compression therapy.

Direction for use

For hygienic hand disinfection apply lotion to hands and keep hands moist for 30 seconds. For surgical hand disinfection apply lotion to hands and forearms and keep them moist for 5 minutes* (*Hygienic expert reports in accordance with DGHM/VAH have certified an application time of 1.5 minutes). For disinfection of skin before simple injections and punctures of peripheral blood vessels apply lotion thoroughly to skin and keep skin moist for 15 seconds. For disinfection before operations and punctures of joints apply lotion thoroughly to skin and keep skin moist for 1 minute. For disinfection of sebaceous skin apply lotion thoroughly to skin and keep skin moist for 10 minutes. For use in cold compression therapy use diluted lotion mixed in equal proportions with water.

Application time

Hygienic hand disinfection	30 sec.
Surgical hand disinfection	1,5 min.
Hand disinfection before injections, punctures	15 sec.
Hand disinfection before operations, punctures of joints, body cavities, hollow organs on un-sebaceous skin	1 min.
on sebaceous skin	10 min.
As bactericidal disinfectant including MRSA, Fungicidal	30 sec.
TB	1 min.
Effectiveness against enveloped viruses including HBV/HIV/HCV/vaccinia/BVDV	30 sec.
Rotaviruses	15 sec.

DGHM/VAH listed

Licensed in line with AMG, licence number: 1599.98.99



Special information

The times given are minimum times. Dependent on additional factors (such as moist skin, unclean skin, risk of the procedure) the application time must be extended. Safe Line Skin Des does not kill spores and is thus unsuitable for the storage of sterile surgical instruments and syringes.

Delivery Forms

Single package	Sales unit	Art.-No.	Ref.-No.
150 ml bottle	1 x 150 ml	780044	09900
250 ml bottle	1 x 250 ml	780045	09901
500 ml bottle	1 x 500 ml	780046	09902
1 litre bottle	1 x 1 L	780047	09903
5 litres container	1 x 5 L	780048	09904

Pharmaceutical expert report

Prof. Dr. Schubert, Frankfurt/Main:

Certification for hand disinfection in accordance with DGHM/VAH norm from 15.04.2010.

PD Dr. Pitten, Gießen:

Certification for hand and skin disinfection in accordance with DGHM/VAH norm from 02.08.2010.

Prof. Kramer/Dr. Rudolph, Greifswald:

Certification for hand disinfection in accordance with DGHM/VAH norm from 26.08.2010.

Dr. Steinmann, virologist, Bremen:

Certification for effectiveness against vaccinia viruses from 20.12.2005.

Certification for effectiveness against BDV viruses from 02.10.2001.

Certification for effectiveness against rotaviruses from 10.06.1992.

Certification for effectiveness against H1N1/H5N1/influenza viruses from 07.04.2010

Warning Notices

Highly inflammable! Keep away from sources of ignition! Keep container tightly closed. In case lotion is spilled certain measures are to be taken immediately in order to avoid fire or explosion. Appropriate measures are such as absorbing lotion, dilution of lotion with water, ventilation of the room, removing of sources of ignition. Keep lotion out of the reach of children. Do not use lotion after expiry date.

Side Effects

Application of lotion can lead to redness of skin and burning skin sensations.

Contraindications

Safe Line Skin Des is unsuitable for disinfection of open skin wounds.

UN 1987